JUN 0 7 2004 JUN 0 7 2004

ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 1 OF 8

OF DT					,					
Troill F10-1449								ete if Know	vn	
U.S. DEPARTMENT OF COMMERCE					Application Number 08/813,829					
PATENT AND TRADEMARK OFFICE				Filin	g Date	Mare	March 6, 1997			
LIST OF INFORMATION CITED BY APPLICANT					First	Named Inventor	Brig	Brigid L.M. Hogan		
(Use as many sheets as necessary)				Grou	up Art Unit	1632	2			
,		•	,,			Examiner Name Joseph Woitach			ch .	
	-:			J.S. PA	CENT	DOCUMENTS	- 300%	- 	- G444075 334	
Examiner's	Cite	Document No.		Date		Name	Class	Subclass	Filing Date (if appropriate	
Initials	No.	5 400 005	445	4/00	1400					
AY	B1 B2	5,166,065	11/2		Williams et al. Green et al.			+ +-		
 	B3	5,334,702 5,605,803	08/1			et al.		 		
 	B4	5,639,618	06/1			et al.	-	 		
 -	B5	5,670,372	09/2		Gay Hogan			 		
	B6	5,690,926	11/19		Hoga		-	 		
 	B7	6,521,671	2/18/			awa et al.	_	 	·	
							30.00	5241 - 22 d 72 m		
Examiner's		Foreign Patent Docu	ıment	Dat		Nam			anslation	
Initials	No.	Country Code-Number-							Yeş/No	
	B8	Kind Code WO 90/03432		04/05/9	an	Evans et al.				
1	B9	WO 97/25412		7/17/97		Anderson et al.				
 	B10	WO 93/17032		9/1993		Hirschmann et	al.			
	1 =	NON-PATIENT DOCUMENTS								
Examiner's	Cite								Place of Publication)	
Initials	No.									
76	B11	Allan et al., 1987, In Perspectives on Mammalian Cell Death, (Potten, C. S. 1Ed.), Oxford University Press, London. pp. 229-258.								
	B12	Allan et al., Spermatogonial apoptosis has three morphologically recognizable phases and								
1 1		shows no circadian rhythm during normal spermatogenesis in the rat. Cell Prolif. 1992 May;25(3):241-50						ell Prolif. 1992		
	<u> </u>									
	B13									
	İ									
			prolite	rative po	otentia	il for prolonged p	eriods	of culture.	Dev. Biol. 227, 271-	
 	B14	278 Anderson, D.J., Gage, F.H., and Weissman, I.L. (2001). Can stem cells cross lineage						roce lineage		
		boundaries? Nat. Med. 7, 393-395.								
	B15	Andrews, P.W. (1998). Teratocarcinomas and human embryology: pluripotent human EC cell lines. Review article. APMIS 106, 158-167								
	B16	Armstrong, L. La					Hole N.	(2000). m	Tert expression	
1 1		correlates with te	lomer	ase activ						
		correlates with telomerase activity during the differentiation of murine embryonic stem cell Mech. Dev. 97, 109-116						·		
	B17	Assady, S., Maor,	G., Ar	nit, M., It	skovitz	-Eldor, J., Skoreck	ci, K.L., a	and Tzuker	man, M. (2001).	
		Insulin production	by hu	man en	bryon	ic stem cells. Dia	betes.	50.		
		http://www.diabetes.org/Diabetes_Rapids/Suheir_Assady_ 06282001.pdf								
B18 Behringer et al., Mullerian-inhibiting substance function during mammalian sex			ian sexual							
	1000	development. Cell. 1994 Nov 4;79(3):415-25. Bellve, Purification, culture, and fractionation of spermatogenic cells. Methods Enzymol.								
	B19	Bellve, Purification		ure, and	fracti	onation of sperm	atogeni	ic cells. M	ethods Enzymol.	
	B20	Bergmann et al.,	Partia						If and horse serum	
<u> </u>		which utilizes the	unrom	DIU LECE	pior s	ystem. J. Cell Bio	JI. 97 (5	paπ 2): 3	94A (1963)."	

Joe Chartees 10/1/0

222303

ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 2 OF 8

DEMARE		
	B21	Bertchtold, A simple method for direct cloning and sequencing cDNA by the use of a single specific oligonucleotide and oligo(dT) in a polymerase chain reaction (PCR). Nucleic Acids Res. 1989 Jan 11;17(1):453.
100	B22	Bhasin et al., Stage dependent expression of inhibin alpha and beta-B subunits during the cycle of the rat seminiferous epithelium. Endocrinology. 1989 Feb; 124(2):987-91991.
	B23	Bicsak et al., Hormonal regulation of inhibin production by cultured Sertoli cells. Mol Cell Endocrinol. 1987 Feb;49(2-3):211-7.
	824	Biesecker and Emerson, "Interleukin-6 is a Component of Human Umbilical Cord Serum and Stimulates Hematopoiesis in Embryonic Stem Cells in vitro" Exp. Hematology 21:774-778 (1993).
	B25	Billington, Biology of the trophoblast. Adv Reprod Physiol. 1971;5:27-66. Review.
	B26	Bogler, O., Wren, D., Barnett, S.C., Land, H. and Noble, M. (1990). Cooperation Between Two Growth Factors Promotes Extended Self-Renewal and Inhibits Differentiations of Oligodendrocyte-type-2 Astrocyte (O-2A) Progenitor Cells. Proc. Natl. Acad. Sci. USA 87, 6368-6372
	B27	Brustle, O., Jones, K.N., Learish, R.D., Karram, K., Choudhary, K., Wiestler, O.D., Duncan, I.D., and McKay, R.D. (1999). Embryonic stem cell-derived gliat precursors: a source of myelinating transplants. Science. 285, 754-756
	B28	Burbank, Douglas W. (October 8, 1992) Characteristic size of relief. Nature 359:483-484.
	B29	Buttery, L.D., Bourne, S., Xynos, J.D., Wood, H., Hughes, F.J., Hughes, S.P., Episkopou, V., and Polak, J.M. (2001). Differentiation of osteoblasts and in vitro bone formation from murine embryonic stem cells. Tissue Eng. 7, 89-99
	B30	Call, L.M., Moore, C.S., Stetten, G., and Gearhart, J.D. (2000). A cre-lox recombination system for the targeted integration of circular yeast artificial chromosomes into embryonic stem cells. Hum. Mol. Genet. 9, 1745-1751
	B31	Cate et al., 1990, In Peptide Growth Factors and Their Receptor, (Sporn et al. Eds.), Springer-Verlag, Berlin, vol. 2, pp. 179-210.
	B32	Cherry, S.R., Biniszkiewicz, D., van Parijs, L., Baltimore, D., and Jaenisch, R. (2000). Retroviral expression in embryonic stem cells and hematopoietic stem cells. Mol. Cell. Biol. 20, 7419-7426
	B33	Chisaka, O. and Capecchi, M.R., "Regionally Restricted Developmental Defects Resulting from Targeted Disruption of the Mouse Homeobox Gene Hox-1.5" Nature 350:473-479 (1991).
	B34	Conlon et al., A primary requirement for nodal in the formation and maintenance of the primitive streak in the mouse. Development. 1994 Jul; 120(7):1919-28.
	B35	Cross et al., Implantation and the placenta: key pieces of the development puzzle. Science. 1994 Dec 2;266(5190):1508-18. Review.
	B36	Dinsmore et al., "High Efficiency Differentiation of Mouse Embryonic Stem Cells into Either Neurons or Skeletal Muscle in vitro" Keystone Symposium (Abstract H111) J. Cell. Biochem. Supplement 18A:177 (1994).
	B37	Donovan & Gerhart, "The end of the beginning for pluripotent stem cells."Nature 414(6859):92-97 (2001)
	B38	Dunnett, S.B., Bjorklund, A., and Lindvall, O. (2001). Cell thera- py in Parkinson's disease—stop or go? Nat Rev. Neurosci. 2, 365-369
	B39	Dzierzak, E. (1999). Embryonic beginnings of definitive hematopoietic stem cells. Ann. N. Y. Acad. Sci. 872, 256-262
	B40	Evans et al., Expression of the helix-loop-helix factor ld during mouse embryonic development. Dev Biol. 1993 Oct;159(2):485-99.
	B41	Fairchild, P.J., Brook, F.A., Gardner, R.L., Graca, L., Strong, V., Tone, Y., Tone, M., Nolan, K.F., and Waldmann, H. (2000). Directed differentiation of dendritic cells from mouse embryonic stem cells. Curr. Biol. 10, 1515-1518

Joe Wolan io/11/00

222303

ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 3 OF 8

	B42	Ffrench-Constant, C., Hollingsworth, A., Heasman, J. and Wylie, C.C. (1991). Response to
1401)	Fibronectin of Mouse Primordial Germ Cells Before, During and After Migration. Development 113, 1365-1373.
1	B43	Finn, The biology of decidual cells. Adv Reprod Physiol. 1971;5:1-26. Review. 5:1-26.
	B44	Fleming et al., "Functional Heterogeneity is Associated with the Cell Cycle Status of Murine
		Hematopoietic Stem Cells" J. Cell Biol. 122:897-902 (1993).
	B45	Frohman et al., Rapid production of full-length cDNAs from rare transcripts: amplification
1		using a single gene-specific oligonucleotide primer. Proc Natl Acad Sci U S A. 1988
I	B46	Dec;85(23):8998-9002 Gallacher, L., Murdoch, B., Wu, D., Karanu, F., Fellows, F., and Bhatia, M. (2000). Identification
	D40	of novel circulating human embryonic blood stem cells. Blood. 96, 1740-1747
	B47	Gardner, D.K. and Schoolcraft, W.B. (1999). Culture and transfer of human blastocysts. Curr.
1	,	Opin. Obstet. Gynecol.
	D40	11, 307-311
1 \ 1	B48	Gavrieli et al., Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. J Cell Biol. 1992 Nov;119(3):493-501.
 	B49	Gearhart, J. (1998). New potential for human embryonic stem cells. Science. 282, 1061-1062
	B50	Godin et al.,(1991). TGFB-1 Inhibits Proliferation and has a Chemotropic Effect on Mouse
		Primordial Germ Cells in Culture. Development 113, 1451-1457.
	B51	Graff et al., Studies with a Xenopus BMP receptor suggest that ventral mesoderm-inducing
		signals override dorsal signals in vivo. Cell. 1994 Oct 7;79(1):169-79.
1 1 1	B52	Green, Mechanism of the pleiotropic effects of the short-ear mutant gene in the mouse. J Exp
 	B53	Zool. 1968 Feb;167(2):129-50. No abstract available. Groves et al., "Repair of Demyelinated Lesions by Transplantation of Purified O-2A Progenitor
	D 33	Cells" Nature 362:453-455 (1993).
	B54	Guenechea, G., Gan, O.I., Dorrell, C., and Dick, J.E. (2001). Distinct classes of human stem cells that differ in prolifera- tive and self-renewal potential. Nat. Immunol. 2, 75-82
	B55	Handel et al., New mutation causing sterility in the mouse. Gamete Res. 1988 Dec;21(4):409-23.
	B56	Haneji et al., "Differential Effects of Epidermal Growth Factor on the Differentiation of Type A Spermatogonia in Adult Mouse Cryptorchid Testes In-Vitro", J. Endocrinol. 128 (3): 383-388 (1991).*
	B57	Harland, The transforming growth factor beta family and induction of the vertebrate mesoderm: bone morphogenetic proteins are ventral inducers. Proc Natl Acad Sci U S A. 1994 Oct 25;91(22):10243-6. Review.
	B58	Harley, C.B., Gearhart, J., Jaenisch, R., Rossant, J., and Thomson, J. (2001). Keystone Symposia. Pluripotent stem cells: biology and applications. Durango, CO
	B59	Hermiston et al. PNAS 90:8866-8870 (1993).
	B60	Hole, N. (1999). Embryonic stem cell-derived haematopoiesis. Cells Tissues Organs. 165, 181-189
	B61	Hong et al., "Production of medakafish chimeras from a stable embryonic stem cell line."PNAS 95(7):3679-3684 (March 31,1998)
	B62	Hooper et al., "HPRT-Deficient (Lesch-Nyhan) Mouse Embryos Derived from Germline Colonization by Cultured Cells" Nature 326:292-295 (1987).
	B63	Hughes, et al., The effects of bone morphogenetic protein-2, -4, and -6 on differentiation of rat osteoblast cells in vitro. Endocrinology. 1995 Jun;136(6):2671-7.
	B64	Itskovitz-Eldor, J., Schuldiner, M., Karsenti, D., Eden, A., Yanuka, O., Amit, M., Soreq, H., and
		Benvenisty, N. (2000). Differentiation of human embryonic stem cells into embry- oid bodies comprising the three embryonic germ layers. Mol. Med. 6, 88-95
	B65	Janatpour et al., The American Society for Cell Biology 34th annual meeting. San Francisco, California, December 10-14, 1994. Abstracts. Mol Biol Cell. 1994 Oct;5 Suppl:453A:2636.

222303

Joe Warler

10/11/06

ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 4 OF 8

· ·
Johnson et al., Two rat homologues of Drosophila achaete-scute specifically expressed in neuronal precursors. Nature. 1990 Aug 30;346(6287):858-61
Jones and Watt, "Separation of Human Epidermal Stem Cells from Transit Amplifying Cells on the Basis of Differences in Integrin Function and Expression" Cell 73:713-723 (1993).
Jones et al., DVR-4 (bone morphogenetic protein-4) as a posterior-ventralizing factor in Xenopus mesoderm induction. Development. 1992 Jun;115(2):639-47.
Jones, G.M., Trounson, A.O., Lolatgis, N., and Wood, C. 1998). Factors affecting the success of human blastocyst development and pregnancy following in vitro fertilization and embryo transfer. Fertil. Steril. 70, 1022-1029
Jones, J.M. and Thomson, J.A. (2000). Human embryonic stem cell technology. Semin. Reprod. Med. 18, 219-223
Kaufman, D.S., Lewis, R.L., Auerbach, R., and Thomson, J.A. (1999). Directed differentiation of human embryonic stem cells into hematopoietic colony forming cells. Blood. 94 (Supplement part 1), 34a
Kehat, I., Kenyagin-Karsenti, D., Druckmann, M., Segev, H., Amit, M., Gepstein, A., Livne, E., Binah, O., Itskovitz-Eldor, J., and Gepstein, L. (2001). Human embryonic stem cells can differentiate into myocytes portraying cardiomyocytic structural and functional properties. J. Clin. Invest. (in press
3 Keller et al., "Hematopoietic Commitment During Embryonic Stem Cell Differentiation in Culture" Mol. Cell. Biol. 13:473-486 (1993).
Khosla, S., Dean, W., Brown, D., Reik, W., and Feil, R. (2001). Culture of preimplantation mouse embryos affects fetal development and the expression of imprinted genes. Biol. Reprod. 64, 918-926
Kim and Smithies, "Recombinant Fragment Assay for Gene Targeting Based on the Polymerase Chain Reaction" Nucleic Acids. Res. 16:8887-8903 (1988).
King et al., BMP5 and the molecular, skeletal, and soft-tissue alterations in short ear mice. Dev Biol. 1994 Nov;166(1):112-22.
Kingsley et al., The mouse short ear skeletal morphogenesis locus is associated with defects in a bone morphogenetic member of the TGF beta superfamily. Cell. 1992 Oct 30;71(3):399-410.
Kingsley, The TGF-beta superfamily: new members, new receptors, and new genetic tests of function in different organisms. Genes Dev. 1994 Jan;8(2):133-46. Review
Kramer, J., Hegert, C., Guan, K., Wobus, A.M., Muller, P.K., and Rohwedel, J. (2000). Embryonic stem cell-derived chondrogenic differentiation in vitro: activation by BMP-2 and BMP-4. Mech. Dev. 92, 193-205
Laughlin, M.J. (2001). Umbilical cord blood for allogeneic transplantation in children and adults. Bone Marrow Transplant. 27, 1-6
Lawson et al., (1992). Clonal Analysis of Cell Fate During Gastrulation and Early Neurulation in the Mouse in CIBA Foundation Symposium 165 Post Implantation Development in the Mouse John Wiley and Sons.
Lee, S.H., Lumelsky, N., Studer, L., Auerbach, J.M., and McKay, R.D. (2000). Efficient generation of midbrain and hindbrain neurons from mouse embryonic stem cells. Nat Biotechnol. 18, 675-679
Lewis, "Embroyonic Stem Cells Debut amid Little Media Attention" The Scientist, 11(19):1(Sept. 29, 1997)
Lillien and Raff, "Differentiation Signals in the CNS: Type-2 Astrocyte Development in vitro as a Model System" Neuron 5:111-119 (1990).
Liu, S., Qu, Y., Stewart, T.J., Howard, M.J., Chakrabortty, S., Holekamp, T.F., and McDonald, J.W. (2000). Embryonic stem cells differentiate into oligodendrocytes and myeli- nate in culture and after spinal cord transplantation. Proc. Natl. Acad. Sci. U. S. A. 97, 6126-6131
Loh et al., Polymerase chain reaction with single-sided specificity: analysis of T cell receptor delta chain. Science. 1989 Jan 13;243(4888):217-20.

222303

Joe Ward 10/11/06

ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 5 OF 8

יט			
	7 U	B87	Lufkin et al., "Disruption of the Hox-1.6 Homeobox Gene Results in Defects in a Region Corresponding to its Rostral Domain of Expression" Cell 66:1105-1119 (1991).
		B88	Lumelsky, N., Blondel, O., Laeng, P., Velasco, I., Ravin, R., and McKay, R. (2001). Differentiation of Embryonic Stem Cells to Insulin-Secreting Structures Similiar to Pancreatic Islets. Science. 292, 1389-1394
		B89	Lyons et al., Organogenesis and pattern formation in the mouse: RNA distribution patterns suggest a role for bone morphogenetic protein-2A (BMP-2A). Development. 1990 Aug;109(4):833-44.
		B90	Mann et al., (1990). Androgenetic Mouse Embryonic Stem Cells are Pluripotent and Cause Skeletal Defects in Chimeras: Implications for Genetic Imprinting. Cell 62, 251-260.
		B91	Manova et al., (1991). Expression of C-Kit Encoded at the W Locus of Mice in Developing Embryonic Germ Cells and Presumptive Melanoblasts. Dev. Biol. 146, 312-324.
		B92	Mansour et al., "Introduction of a LacZ Reporter gene into the Mouse int-2 Locus by Homologous Recombination" Proc. Natl. Acad. Sci. USA 87:7688-7692 (1990).
		B93	Massague et al., The TGF-beta family and its composite receptors. Trends Cell Biol. 1994 May;4(5):172-8.
		B94	Matzuk et al., Alpha-inhibin is a tumour-suppressor gene with gonadal specificity in mice. Nature. 1992 Nov 26;360(6402):313-9.
		B95	McDonald, J.W., Liu, X.Z., Qu, Y., Liu, S., Mickey, S.K., Turetsky, D., Gottlieb, D.I., and Choi, D.W. (1999). Transplanted embryonic stem cells survive, differentiate and promote recovery in injured rat spinal cord. Nat. Med. 5, 1410-1412
		B96 \	McKay, R., "The Origins of Cellular Diversity in the Mammalian Central Nervous System" Cell 58:815-821 (1989).
		B97	McMahon et al., "The Midbrain-Hindbrain Phenotype of Wnt-1.sup Wnt-1.sup Mice Results from Stepwise Deletion of engrailed-Expressing Cells by 9.5 Days Postcoitum" Cell 69:581-595 (1992).
		B98	Meunier et al., Gonadal and extragonadal expression of inhibin alpha, beta A, and beta B subunits in various tissues predicts diverse functions. Proc Natl Acad Sci U S A. 1988 Jan;85(1):247-51
		/ B99	Meyer et al., "Serum basic fibroblast growth factor in men with and without prostate carcinoma", Cancer 76 (11): 2304-11 (1995).*
	ĺ	B100	Momma, S., Johansson, C.B., and Frisen, J. (2000). Get to know your stem cells. Curr. Opin. Neurobiol. 10, 45-49.
		B101	Morrison, S.J. (2001). Neuronal differentiation: Proneural genes inhibit gliogenesis. Curr. Biol. 11, R349-R351.
		B102	Murray et al., Bone morphogenetic protein inhibits differentiation and affects expression of helix-loop-helix regulatory molecules in myoblastic cells. J Cell Biochem. 1993 Sep;53(1):51-60.
		B103	Nishikawa, S.I., Nishikawa, S., Hirashima, M., Matsuyoshi, N., and Kodama, H. (1998). Progressive lineage analysis by cell sorting and culture identifies FLK1(+)VE-cadherin(+) cells at a diverging point of endothelial and hemopoietic lineages. Development. 125, 1747-1757
		B104	Noble, M., Murray, K., Stroobant, P., Waterfield, M.D., and Riddle, P. (1988). Platelet-derived growth factor promotes division and motility and inhibits premature differentiation of the oligodendrocyte/type-2 astrocyte progenitor cell. Nature 333, 560-562.
		B105	Odorico, J.S., Kaufman, D.S., and Thomson, J.A. (2001). Multilineage Differentiation from Human Embryonic Stem Cell Lines. Stem Cells. 19, 193-204
	6	B106	Ott et al., "Use of Recombinant Embryonic Stem Cells to Isolate Neural Stem Cells" Keystone Symposium (Abstract H222) J. Cell. Biochem. Supplement 18A:187 (1994).
		B107	Ozkaynak et al., Osteogenic protein-2. A new member of the transforming growth factor-beta superfamily expressed early in embryogenesis. J Biol Chem. 1992 Dec 15;267(35):25220-7.

222303

Joe Worter

10/11/06



ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 6 OF 8

Ju	B108	Panicker, M. and Rao, M. (2001). Stem cells and neuro- genesis. Marshak, D.R., Gardner, D.K., and Gottlieb, D. eds. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press). 399-438.
f	B109	Parvinen et al., "In-Vitro Stimulation of Stage-Specific DNA Synthesis in Rat Seminiferous Tublule Segments by Interleukin 1-alpha" Endocrinology 129 (3): 1614-20 (1991).*
	B110	Patterson, P., "Control of Cell Fate in a Vertebrate Neurogenic Lineage" Cell 62:1035-1038 (1990).
	B111	Pedersen, R.A. (1999). Embryonic stem cells for medicine. Sci. Am. 280, 68-73
	B112	Pera, M.F., Reubinoff, B., and Trounson, A. (2000). Human embryonic stem cells. J. Cell Sci. 113 (Pt 1), 5-10
	B113	Perkins, A.C. (1998). Enrichment of blood from embryonic stem cells in vitro. Reprod. Fertil. Dev. 10, 563-572
	B114	Piedrahita et al., "Generation of transgenic porcine chimeras using primordial germ cell-derived colonies," Biological Reproduction 58(5):1321-9 (1998) Abstract Only
	B115	Porteus et al., "Isolation and Characterization of a Novel cDNA Clone Encoding a Homeodomain That is Developmentally Regulated in the Ventral Forebrain" Neuron 7:221-229 (1991).
1	B116	Price et al., "A Mouse Gene Related to distal-less Shows a Restricted Expression in the Developing Forebrain" Nature 351:748-751 (1991).
	B117	Rathjen, P.D., Lake, J., Whyatt, L.M., Bettess, M.D., and Rathjen, J. (1998). Properties and uses of embryonic stem cells: prospects for application to human biology and gene therapy. Reprod. Fertil. Dev. 10, 31-47
	B118	Reddy et al. Fluorescence-activated sorting of totipotent embryonic stem cells expressing developmentally regulated lacZ fusion genes. Proc Natl Acad Sci U S A. 1992 Aug 1;89(15):6721-5.
	B119	
	B120	Reubinoff, B.E., Pera, M.F., Fong, C.Y., Trounson, A., and Bongso, A. (2000). Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. Nat. Biotechnol. 399-404
	B121	Robey, P.G. (2000). Stem cells near the century mark. J. Clin. Invest. 105, 1489-1491
	B122	Roelink, H. and Nuse, R., "Expression of Two Members of the Wnt Family During Mouse Development-Restricted Temporal and Spatial Patterns in the Developing Neural Tube" Genes Dev. 5:381-388 (1991).
	B123	Rossi et al., "Follicle-Stimulating Hormone Induction of Steel Factor SLF mRNA in Mouse Sertoli Cells and Stimulation of DNA Synthesis in Spermatogonia by Soluble SLF", Dev. Biol. 155 (1): 68-74 (1993).*
	B124	Rowe et al., Maps from two interspecific backcross DNA panels available as a community genetic mapping resource. Mamm Genome. 1994 May;5(5):253-74. Erratum in: Mamm Genome 1994 Jul;5(7):463.
	B125	Rudnicki et al., Inactivation of MyoD in mice leads to up-regulation of the myogenic HLH gene Myf-5 and results in apparently normal muscle development. Cell. 1992 Oct 30;71(3):383-90
	B126	Russell et al. 1990, In Histological and Histopathological Evaluation of the Testis, Cache River Press, Clearwater, Fl., pp. 119-161.
	B127	Sandberg et al., "Reviews of chromosome studies in urological tumors. III. Cytogenetics and genes in testicular tumors, "J. Urology 155(5):1531-56 (1996) Abstract
	B128	Sanger et al., DNA sequencing with chain-terminating inhibitors. Proc Natl Acad Sci U S A. 1977 Dec;74(12):5463-7. No abstract available.
	B129	Schmitt et al., "Hematopoietic Development of Embryonic Stem Cells in vitro: Cytokine and Receptor Gene Expression" Genes and Develop. 5:728-740 (1991).

222303

Joe Warted 10/11/06

ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 7 OF 8

_	-82		
RADE	MIREST.	B130	Schuldiner, M., Yanuka, O., Itskovitz-Eldor, J., Melton, D., and Benvenisty, N. (2000). Effects of eight growth factors on the differentiation of cells derived from human embryonic stem cells.
-	711	D424	Proc. Natl. Acad. Sci. U. S. A. 97, 11307-11312
ı	7	B131	Shamblott, M.J., Axelman, J., Littlefield, J.W., Blumenthal, P.D., Huggins, G.R., Cui, Y., Cheng, L.,
	1		and Gearhart, J.D. (2001). Human embryonic germ cell derivatives express a broad range of
			develpmentally distinct markers and proliferate extensively in vitro. Proc. Natl. Acad. Sci. U. S. A. 98, 113-118
		B132	Shamblott, M.J., Axelman, J., Wang, S., Bugg, E.M., Littlefield, J.W., Donovan, P.J., Blumenthal,
l	- 1		P.D., Huggins, G.R., and Gearhart, J.D. (1998). Derivation of pluripotent stem cells from cultured
			human primordial germ cells. Proc. Natl. Acad. Sci. U. S. A. 95, 13726-13731
1	- 1	B133	Shen et al. Embryonic stem cells stably transfected with mRAR beta 2-lacZ exhibit specific
L			expression in chimeric embryos. Int J Dev Biol. 1992 Dec;36(4):465-76.
		B134	Simeone et al., "Nested Expression Domains of Four Homeobox Genes in Developing Rostral Brain" Nature 358:687-690 (1992).
		B135	Simeone et al., "Two Vertebrate Homeobox Genes Related to the Drosophila empty spiracles Gene are Expressed in the Embryonic Cerebral Cortex" EMBO J. 11(7):2541-2550 (1992)
		B136	Smith, A.G. (2001). Origins and properties of mouse embryonic stem cells. Annu. Rev. Cell. Dev. Biol 17:435-462
		B137	Snodgrass et al., "Embryonic Stem Cells and in vitro Hematopoiesis" J. Cell. Biochem. 49:225-230 (1992).
		B138	Steinberger et al., Secretion of an FSH-inhibiting factor by cultured Sertoli cells. Endocrinology. 1976 Sep;99(3):918-21921.
		B139	Storm et al., Limb alterations in brachypodism mice due to mutations in a new member of the TGF beta-superfamily. Nature. 1994 Apr 14;368(6472):639-43.
		B140	Thomson, J.A. and Marshall, V.S. (1998). Primate embryonic stem cells. Curr. Top. Dev. Biol. 38, 133-165
		B141	Thomson, J.A., Itskovitz-Eldor, J., Shapiro, S.S., Waknitz, M.A., Swiergiel, J.J., Marshall, V.S., and Jones, J.M. (1998). Embryonic stem cell lines derived from human blastocysts. Science. 282,
\perp			1145-1147
		B142	Trounson, A.O., Anderiesz, C., and Jones, G. (2001). Maturation of human oocytes in vitro and their develop- mental competence. Reproduction. 121, 51-75
	1	B143	Tsai, M., Wedemeyer, J., Ganiatsas, S., Tam, S.Y., Zon, L.I., and Galli, S.J. (2000). In vivo
			immunological function of mast cells derived from embryonic stem cells: an approach for the
			rapid analysis of even embryonic lethal mutations in adult mice in vivo. Proc. Natt. Acad. Sci. U.
-	- -	B144	S. A. 97, 9186-9190 Tsokos, G.C. and Nepom, G.T. (2000). Gene therapy in the treatment of autoimmune
	/ /	5144	diseases. J. Clin. Invest 106,
L	/		181-183
	$\neg T$	B145	Vassalli et al., Activin/inhibin beta B subunit gene disruption leads to defects in eyelid
-		D140	development and female reproduction. Genes Dev. 1994 Feb 15;8(4):414-27
		B146	Weissman, I.L. (2000). Stem cells: units of development, units of regeneration, and units in evolution. Cell. 100,157-168
		B147	Wiles, M.V., Vauti, F., Otte, J., Fuchtbauer, E.M., Ruiz, P., Fuchtbauer, A., Amold, H.H., Lehrach.
1	1		H., Metz, T., von Melchner, H., and Wurst, W. (2000). Establishment of a gene-trap sequence
L			tag library to generate mutant mice from embryonic stem cells. Nat. Genet. 24, 13-14.
		B148	Wolswijk and Noble, "Identification of an Adult-Specific Glial Progenitor Cell" Development 105:387-400 (1989).
		B149	Wright et al., Isolation of cyclic protein-2 from rat seminiferous tubule fluid and Sertoli cell culture medium. Biol Reprod. 1986 Oct;35(3):761-72.
	7	B150	Wright et al., 1993, In Cell and Molecular Biology of the Testis, (Eds.: Desjardins et al.),
	<u> </u>		pp.377-399, Oxford Univ. Press NY.

222303

Joe (Voilage) 10/11/06

ATTORNEY DOCKET NO. 16016.0005US APPLICATION NO. 08/813,829 SHEET 8 OF 8

Αl	JE WILL				
	Ap	B151	Yagi et al., Homologous recombination at c-fyn locus of mouse embryonic stem cells with use of diphtheria toxin A-fragment gene in negative selection. Proc Natl Acad Sci U S A. 1990 Dec;87(24):9918-22.		
		B152	Yamashita, J., Itoh, H., Hirashima, M., Ogawa, M., Nishikawa, S., Yurugi, T., Naito, M., Nakao, K., and Nishikawa, S. (2000). Flk1-positive cells derived from embryonic stem cells serve as vascular progenitors. Nature. 408, 92-96		
		B153	Zhao et al., Cartilage homeoprotein 1, a homeoprotein selectively expressed in chondrocytes. Proc Natl Acad Sci U S A. 1993 Sep 15;90(18):8633-7.		
		B154	Zhou et al., Nodal is a novel TGF-beta-like gene expressed in the mouse node during gastrulation. Nature. 1993 Feb 11;361(6412):543-7.		
	-	B155	Materials from National Institutes of Health website called <u>Stem Cells: Scientific Progress and Future Research Directions</u> (June 2001)		
	Examiner	Signatu	re: Prof (1416) Date Considered: 10/11/06		
	EXAMINER: Initial if reference considered, whether or not citation is in conformance with MRER 600: Draw line				